

RESEARCH ARTICLE

Determination of the levels of heavy metals and physicochemical properties of borehole water within selected mining sites in Ebonyi State, Nigeria

David Okechukwu Okeke^{1*} Jonathan Chinenye Ifemeje² Victor Chukwuemeka Eze³¹ Department of Applied Biochemistry, Nnamdi Azikiwe University, Awka, Anambra state, Nigeria² Department of Biochemistry, Chukwuemeka Odumegwu Ojukwu University, Uli, Anambra State, Nigeria³ Department of Pure and Industrial Chemistry, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria

Correspondence to: David Okechukwu Okeke, Department of Applied Biochemistry, Nnamdi Azikiwe University, P.M.B. 5025, Awka, Anambra state, Nigeria; E-mail: do.okeke@unizik.edu.ng

Received: August 28, 2021;**Accepted:** February 17, 2022;**Published:** February 25, 2022.

Citation: Okeke DO, Ifemeje JC and Eze VC. Determination of the levels of heavy metals and physicochemical properties of borehole water within selected mining sites in Ebonyi State, Nigeria. *Health Environ*, 2022, 3(1): 162-168.
<https://doi.org/10.25082/HE.2022.01.003>

Copyright: © 2022 David Okechukwu Okeke *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by-nc/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Abstract: The level of heavy metals (Pb, Cd, As, Cr, Ni, Co, Zn) and physicochemical parameters in borehole water sources within selected mining sites in Ebonyi State, Nigeria were determined using FS240AA Atomic Absorption Spectrophotometer (AAS) according to the method of American Public Health Association (APHA). Water samples were collected from Enyigba mining site, Ikwo mining site, Ameri Amekamining site, Izza mining site, Mkpume Akwatakwa mining site and Mpume Akwaokuku mining sites. Composite water samples were collected within the mining locales and 500 metres away from the mining locations using two liters plastic jars. Similarly, the control samples were taken from Abakaliki (about 15 km away from mining area) where there was no evidence of mining activity. A total of 14 sub-samples were collected for this study. Generally, the levels of heavy metals in the borehole water samples were higher than the values recommended by the World Health Organization (WHO). This however suggests a possible mobility of the metals from mining sites into the groundwater sources through leaching. The findings from this study have revealed the need to purify borehole water within mining areas prior to industrial and domestic use.

Keywords: heavy metals, physicochemical, borehole, mining site, Ebonyi State

1 Introduction

Water is an inorganic, transparent, tasteless, odorless, and nearly colorless chemical substance, which is vital for sustaining global socio-economic development and all known forms of life [1]. It has been established that water covers approximately 70.9% of the Earth's surface, mostly in seas and oceans. Water can be categorized into two main sources viz groundwater sources and surface water sources. Groundwaters are beneath the land surface and include boreholes, springs and wells, while surface waters are above the land surface and include streams, creeks, wetlands and reservoirs [2]. Small portions of water occur as groundwater (1.7%), in the glaciers and the ice caps of Antarctica and Greenland (1.7%), and in the air as vapor, clouds (consisting of ice and liquid water suspended in air), and precipitation (0.001%) [3]. Heavy metals often contaminate rivers thereby making them unfit for domestic, agricultural and industrial purposes [4,5]. Due to the unique characteristics of heavy metals (high toxicity at low concentrations, poor biodegradability and bioaccumulation), they pose a serious health threat to humans when ingested [5,6]. In Nigeria, borehole water sources are considered the main source of potable drinking water due to the collapse of public pipe-borne water scheme [7-9]. The borehole water resources are usually accessed through the drilling of boreholes and with the help of pumping machines are subsequently pumped out through pipes and then stored in tanks for domestic and industrial use [10]. According to Ibe *et al.* [10], majority of the boreholes within urban and sub-urban areas are constructed close to waste dumpsites, latrines, vicinity of metal scrap yards, mining sites etc., hence rendering the groundwater sources vulnerable to contamination. There are extensive reports that groundwater resources in developing countries like Nigeria are appreciably polluted due to anthropogenic activities such as mining [11,12]. A study conducted by Duru *et al.* [13] attributed the poor water quality in groundwater samples collected in Orji to high cadmium concentration caused by anthropogenic activities. This study is very important owing to the fact that residents within the mining sites depend solely on borehole water resources for their household water requirements; hence the need to regularly monitor borehole water resources within the mining sites to ascertain the quality of the water. Also, this study is vital owing to the paucity of data on the level of heavy metals in borehole

water resources within the study area. The present study, therefore, is aimed at ascertaining the level of heavy metals in borehole as well as the physicochemical properties of the water. The report of this study would, therefore, create the necessary awareness and consciousness amongst the inhabitants, town planners, and other relevant government agencies that will lead to the planning towards the supply and development of future sustainable water schemes in the area.

2 Materials and Methods

2.1 Study area

The study was conducted in six mining sites, namely: Enyigba, Ikwo, Ameri-Ameka, Izza, Mkpume-Akwatakwa, Mpume-Akwaokuku, and the control site (Abakaliki town). The Enyigba mining site is one of the significant communities where mining is vigorously done. It is situated in the south of Abakaliki; 14 km south of Abakaliki town of Ebonyi State [14–16]. Previous investigations at the examination area zeroed in on the effect of heavy metal tainting on deserted mine pits, underground water sources, waterways, streams and counterfeit lakes [11, 17]. Ikwo is located on the eastern part of the state. The territory is plentiful in mineral assets and progenitors of the present occupants created bronze projecting methods in excess of 500 years prior. Ameri-Ameka is situated in the south of Abakaliki town. It comprises of cretaceous dregs of the Asu stream bunch, predominantly shales, silty shales, limestone and volcanic rocks. They are one of Africa’s most notable lead-zinc mineralized region [11, 18]. The Izza south mining site is situated in the south of Abakaliki town. There are various kinds of soil around there, specifically: lateritic mud soil, sandy soil with mudstone and clayey shaly soil. The soil is gotten predominantly from the old sedimentary rocks. The Mkpume-Akwaokuku and Akwatakwa mining sites are situated in the north of Abakaliki town. The vegetation is overwhelmed by grasses, bushes and trees such as palm trees, coconut, mango and orange trees [11, 19, 20]. (see in Table 1, Figure 1)

Table 1 Sampling site and their coordinates

Sampling sites	Coordinates
Enyigba mining site	N: 06° 11.642' E: 008° 08.380' 263.4ft
Ikwo mining site	N: 06° 10.590' E: 008° 07.438' 196.8ft
Ameri Amekaming site	N: 06° 11.042' E: 008° 06.110' 149.3ft
Izza mining site	N: 06° 09.929' E: 008° 06.625' 161.9ft
Mkpume Akwatakwa mining site	N: 06° 23.588' E: 008° 09.784' 192.8ft
Mpume Akwaokuku mining site	N: 06° 30.268' E: 008° 16.352' 226.6ft
Control (Abakaliki town)	N: 06° 19.163' E: 008° 06.559' 142.8ft




Figure 1 Map of Ebonyi State showing the sampling sites

2.2 Samples collection and preparation

In this study, six sampling locations were arbitrarily selected for borehole water sampling using a global positioning system (Model, GARMIN etrex 20). The sampling locations are: Enyigba mining site, Ikwo mining site, Ameri-Ameka mining site, Izza mining site, Mkpume-Akwatakwa mining site and Mpume-Akwaokuku mining site. Composite water samples were collected using two liters plastic jars within the mining locales and 500 metres away from the mining locations [11]. Prior to filling the jars, each was washed thrice with the water from the borehole before sampling. One ml of Nitric corrosive was added to 1 litre of the plastic can which was utilized for cation and anions investigation. The control samples were taken from Abakaliki (about 15 km away from the study area) where there was no evidence of mining activity. A total of 14 sub-samples were collected for this study. The samples were promptly moved to Springboard Research Laboratory, Awka, for analysis.

2.3 Assessment of physicochemical properties of the water samples

The collected water samples were analyzed according to the method prescribed by the American Public Health Association, APHA [21,22]. The pH and electrical conductivity ($\mu\text{s}/\text{cm}$) of the samples were measured *in situ* using pH meter and conductivity meter respectively [22]. Total dissolved solvent (TDS) was determined using APHA [22] while temperature and turbidity were determined using APHA [21]. Hardness (NTU) was conducted by EDTA titrimetric method [22]. Chloride (Cl^-) and fluoride (F^-) were estimated using the Argentometric method [21]. Nitrate (NO_3^-) was estimated using the procedures given by APHA [22].

2.4 Methods of heavy metal analysis of water samples

Heavy metal analysis was conducted using FS240AA Atomic Absorption Spectrophotometer according to the method of American Public Health Association [21]. The samples were thoroughly mixed by shaking, and 100ml of it was transferred into a glass beaker of 250 ml, to which 5ml of concentrated nitric acid was added and heated to boil till the volume is reduced to about 15-20ml. 5ml of concentrated nitric acid was added periodically until all the residues were completely dissolved. The mixture was cooled, transferred and made up to 100 ml using metal free distilled water. The sample was aspirated into the oxidising air-acetylene flame. When the aqueous sample was aspirated, the sensitivity for 1% absorption was observed. The instrument setting and operational conditions were done in accordance with the manufacturers' specifications. Following the same protocol as the samples, standard reference material (SRM 2783) filter from National Institute of Standards and Technology (NIST) was analysed for elemental concentrations and compared with their certified values to validate the analyses and results were found within $\pm 5\%$ of the certified values [19, 23, 24].

2.5 Data analysis

The data analysis for the data reported in this study was done using Ms-Excel.

3 Results and discussions

3.1 Physicochemical parameters in water samples

The results obtained in Izza mining site indicated that all the physicochemical properties of the water samples collected within the mining sites varied significantly from the water samples collected 500 m away from mining site. For Ikwo mining sites, nitrate and hardness did not vary from samples collected 500 m from mining sites. In Akwatakwa mining sites, pH and Hardness did not vary significantly from samples collected 500 m from mining site ($p > 0.05$). From samples collected from Ameri ameka mining site, no variation was observed between pH and TDS in comparison with the control sample. From Mpume akwaokuku, there was no variation between hardness and fluoride in comparison with samples collected 500 m from the mining site. It is worth noting that pH of the borehole water samples were acidic (5.25 – 6.37). Eze et al. [23] stated that daily ingestion of water with low pH could lead to peptic ulcer. The values reported for chloride and fluoride were below the WHO [30] and the NSDWQ acceptable limits. However, the presence of Cl^- and F^- in the borehole water samples are a sign of pollution due

to human activities [23]. The result of the physicochemical parameters assessed in the water samples collected around the mining sites and 500 m from mining site are shown in Table 2.

Furthermore, the low nitrate concentrations recorded in the borehole water samples indicates a low possibility of the development of methaemoglobinaemia in children (infants). In this study, the results reported for the physicochemical parameters were higher than that reported by Duru et al. [13] and Ibe et al. [25].

Table 2 Physicochemical properties of water samples from the selected mining sites

Parameters	Enyigba	Enyigba (500 m)	Izza south	Izza south (500 m)	Ikwo	Ikwo (500 m)	Akwatakwa	Akwatakwa (500 m)	Ameri Ameka	Ameri Ameka (500 m)	Mpume-Akwaokuku	Mpume-Akwaokuku (500 m)	Control
pH	5.39	6.18	5.62	6.37	5.64	5.90	5.51	5.25	5.94	5.85	5.63	5.74	6.44
Temperature (°C)	28.40	29.37	29.20	27.73	27.40	28.53	28.3	26.57	27.4	28.27	18.90	29.33	28.40
Conductivity (us/cm)	190.963	68.83	170.04	66.27	83.72	64.33	62.65	57.30	52.393	86.63	66.05	86.27	45.53
Turbidity (NTU)	12.70	4.37	10.67	5.80	12.83	3.63	16.4	3.57	11.5	3.6	14.80	4.43	2.00
Chloride (mg/l)	174.80	113.43	159.97	93.10	163.96	104.93	141.97	71.98	133.36	82.977	123.43	104.79	122.56
Fluoride (mg/l)	1.25	0.06	1.34	0.16	1.063	0.16	1.82	0.44	1.55	0.347	1.34	0.17	0.25
Nitrate (mg/l)	9.93	8.69	5.28	4.54	6.676	6.37	5.55	3.48	5.51	4.123	11.57	3.47	4.41
TDS (mg/l)	123.33	88.67	123.33	105.33	79.33	65.67	104.67	55.00	73.63	92.00	91.33	81.67	93.00
Hardness (mg/l)	144.33	113.67	133.33	106.00	89.00	126.33	72.00	143.33	79.00	106.66	78.33	107.00	67.72

3.2 Heavy metal concentration in water samples

The result of heavy metal concentration (Table 3) showed a higher level of lead in all the mining sites with Enyigba mining site having the highest concentration of 4.71 mg/l which is followed by Akwatakwa mining site (3.64 mg/l), Izza south (3.40 mg/l), Ikwo (2.75 mg/l), Mkpume-Akwaokuku (2.62 mg/l) and AmeriAmeka (1.56 mg/l). Lower concentrations of lead were detected in the water samples collected 500 m from the mining site with Akwatakwa having the highest concentration of 0.24 mg/l. Generally, low concentration of the heavy metals were detected in both the control samples and the samples collected from the mining sites.

In addition, most of the heavy metals detected recorded higher concentration in samples from the mining site when compared to the samples collected 500 m away from mining site and control sample; except for cobalt which was had higher concentrations in samples collected 500 m away from mining site.

Table 3 Heavy metal analysis of water samples within the selected mining sites

Heavy metals	Enyigba	Enyigba (500 m)	Izza south	Izza south (500 m)	Ikwo	Ikwo (500 m)	Akwatakwa	Akwatakwa (500 m)	Ameri Ameka	Ameri Ameka (500 m)	Mpume-Akwaokuku	Mpume-Akwaokuku (500 m)	Control
Lead (mg/l)	4.713	0.078	3.400	0.065	2.754	0.061	3.635	0.236	1.560	0.055	2.619	0.065	0.122
Cadmium (mg/l)	0.408	0.00	0.257	0.00	0.355	0.023	0.201	0.017	0.156	0.015	0.133	0.025	0.055
Arsenic (mg/l)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Chromium (mg/l)	0.0377	0.00	0.038	0.00	0.022	0.00	0.032	0.00	0.161	0.00	0.035	0.00	0.00
Nickel (mg/l)	0.358	0.02	0.13	0.014	0.28	0.021	0.108	0.025	0.145	0.023	0.351	0.027	0.034
Cobalt (mg/l)	0.00	0.022	0.00	0.017	0.00	0.027	0.00	0.014	0.00	0.027	0.00	0.019	0.00
Zn (mg/l)	6.0135	6.249	4.875	4.361	5.686	6.889	7.087	4.327	6.8	3.722	8.615	2.645	2.654
Sum	11.5302	6.369	8.7	4.457	9.097	7.021	11.063	4.619	8.822	3.842	11.753	2.781	2.865

The result of the current study is comparable to the study of groundwater samples near waste dumpsite conducted by Nnaji et al. [2]. Eze et al. [23] stated that lead causes diseases of the kidney, fetal brain, and damage to the circulatory and nervous system. However, Calo and Parise [26] in their related study reported concentrations of Cd, As, Co, Ni and Pb lower than the results of the current study.

3.3 Assessment of pollution status of water samples

3.3.1 Contamination factors of heavy metals in water

As shown in Table 4, lead (Pb) recorded the highest contamination factor in all the mining sites; cadmium was higher in Enyigba (7.41) and Ikwo (6.45) mining sites, nickel was higher in Enyigba (10.53), Ikwo (8.24) and Mpume akwaokuku (10.32) mining site. However, the results showed a significantly higher contamination around the mining site in comparison with samples 500 m away from the mining site. A moderate to low contamination factor were detected in water samples 500 m away from mining site. This was in agreement with the study conducted by Wongsasuluk et al. [27] and Mishra et al. [28].

3.3.2 Pollution index

The result of pollution indices of heavy metals in water samples collected within the studied mining site (Table 5) showed a higher pollution index for samples collected within mining site in comparison with samples collected 500 m away from the mining site. Moreover, all the samples

Table 4 Contamination factors of heavy metals in water samples from the studied mining sites Vs. (500 m away)

Parameters	Enyigba mining	Izza south mining	Ikwo mining	Akwatakwa mining	Ameri ameka mining	Mpume-Akwaokuko mining
Lead (mg/l)	38.63* (0.64)	27.87* (0.53)	22.57* (0.50)	29.80* (1.93)	12.79* (0.45)	21.47* (0.53)
Cadmium (mg/l)	7.41* (0.00)	4.67 (0.00)	6.45* (0.42)	3.65 (0.31)	2.84 (0.27)	2.42 (0.45)
Arsenic (mg/l)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Chromium (mg/l)	3.77 (0.00)	3.80 (0.00)	2.20 (0.00)	3.20 (0.00)	16.10* (0.00)	3.50 (0.00)
Nickel (mg/l)	10.53* (0.59)	3.82 (0.41)	8.24* (0.62)	3.18 (0.74)	4.26 (0.68)	10.32* (0.79)
Cobalt (mg/l)	0.00 (2.20)	0.00 (1.70)	0.00 (2.70)	0.00 (1.40)	0.00 (2.70)	0.00 (1.90)
Zinc (mg/l)	2.26 (2.35)	1.84 (1.64)	0.00 (2.60)	2.67 (1.63)	2.56 (1.40)	3.25 (1.00)

Note: low: $Cf < 1$, moderate: $1 \geq Cf \leq 3$, Moderately high: $3 > Cf \leq 6$, very high: $Cf > 6$. * indicates high contamination within the study area

collected recorded a high pollution level indicating the influence of anthropogenic activities within the study area. Enyigba mining site recorded a higher pollution index (112.05) around the mining site; while Akwatakwa recorded the highest pollution load index (7.21) 500 m away from the mining site. Generally, the borehole water in the study area was polluted and this could result to adverse health effects on humans who consume the water [23, 29, 30].

Table 5 Pollution Index of heavy metals in water samples from the studied mining sites

Study area	Mining sites	500 m away
Enyigba	112.05	6.31
Izza South	100.013	4.75
Ikwo	65.74	6.84
Akwatakwa	73.98	7.21
Ameri ameka	61.36	6.03
Mpumeakwaokuku	62.59	5.37

Note: low: $PI \leq 1$, middle: $(1 < PI \leq 3)$ or high: $(PI > 3)$

4 Conclusion

This research has revealed that the water samples are acidic (low pH), with a moderate electrical conductivity, total dissolved solid, nitrate, chloride, fluoride and hardness. The ambient values reported are a clear indication of the influence of mining activities within the study area. The local populations who consume water from the boreholes are hereby advised to purify the water before usage. Generally, the heavy metal levels in the water samples from mining communities were higher within the mining sites than that of the samples collected 500 m away from the mining site. This could be associated with mining activities and the high level of air pollution that typically characterizes mining sites.

References

- [1] Water Q & A: Why is water the “universal solvent”? www.usgs.gov. (U.S. Department of the Interior). Retrieved 15 January 2021.
- [2] Nnaji AO, Idam JO and Njoku RE. Assessment of variations in the concentration of heavy metals in groundwater within oil-producing communities of Rivers State Nigeria. *Journal of Public Health & Environment Pollution*, 2016, **1**(1): 1-13.
- [3] Isiuku BO and Enyoh CE. Water pollution by heavy metal and organic pollutants: Brief review of sources, effects, and progress on remediation with aquatic plants. *Analytical Methods in Environmental Chemistry Journal*, 2019, **2**(3): 5-38.
<https://doi.org/10.24200/amecj.v2.i03.66>
- [4] Pazand K, Khosravi D, Ghaderi MR, *et al.* Hydrogeochemistry and lead contamination of groundwater in the north part of Esfahan province, Iran. *Journal of Water & Health*, 2018, **16**(4): 622-634.
<https://doi.org/10.2166/wh.2018.034>
- [5] Kumar M, Ramanathan AL, Tripathi R, *et al.* A study of trace element contamination using multivariate statistical techniques and health risk assessment in groundwater of Chhaprola Industrial Area, Gautam Buddha Nagar, Uttar Pradesh, India. *Chemosphere*, 2017, **166**: 135-145.
<https://doi.org/10.1016/j.chemosphere.2016.09.086>
- [6] Barzegar R, Asghari M, Soltani EF, *et al.* Heavy metal (loid)s in the groundwater of Shabestar area (NW Iran): source identification and health risk assessment. *Exposure and Health*, 2019, **11**: 251-265.
<https://doi.org/10.1007/s12403-017-0267-5>
- [7] Obot EE and Abasifreke EA. Spatial distribution and variability of groundwater quality in the state capital and contiguous Local Government Areas under urbanization expansion. *Journal of Water Resources Planning & Management*, 2014, **2**(1): 1-9.

- [8] Nnaji AO, Idam JO and Njoku RE. Assessment of variations in the concentration of heavy metals in groundwater within oil-producing communities of Rivers State Nigeria. *Journal of Public Health & Environment Pollution*, 2016, **1**(1): 1-13.
- [9] Li F, Qiu Z, Zhang J, *et al.* Investigation, pollution mapping and simulative leakage health risk assessment for heavy metals and metalloids in groundwater from a typical brownfield, middle China. *International Journal of Environmental Research and Public Health*, 2017, **14**(7): 768. <https://doi.org/10.3390/ijerph14070768>
- [10] Ibe FC, Enyoh CE, Opara AI, *et al.* Evaluation of pollution status of groundwater resources of parts of Owerri metropolis and environs, Southeastern Nigeria, using health risk and contamination models. *International Journal of Energy and Water Resources*, 2020, **4**: 357-374. <https://doi.org/10.1007/s42108-020-00071-8>
- [11] Okeke DO, Ifemeje JC and Eze VC. Levels Of Heavy Metals In Soils and Food Crops Cultivated Within Selected Mining Sites In Ebonyi State, Nigeria. *Health and Environment*, **2**(1): 84-95. <https://doi.org/10.25082/HE.2021.01.003>
- [12] Obasi IA, Nnachi EE, Igwe OE, *et al.* Evaluation of pollution status of heavy metals in the groundwater system around open dumpsites in Abakaliki urban, Southeastern Nigeria. *African Journal of Environment Science & Technology*, 2015, **9**(7): 600-609. <https://doi.org/10.5897/AJEST2015.1780>
- [13] Duru CE, Okoro IP and Enyoh CE. Quality assessment of borehole water within Orji mechanic village using pollution and contamination models. *Int J Chem Mat Environ Res*, 2017, **4**(3): 123-130.
- [14] Antoniadis V, Shaheen SM, Boersch J, *et al.* Bioavailability and Risk Assessment of Potentially Toxic Elements in Garden Edible Vegetables and Soils around a Highly Contaminated Former Mining Area in Germany. *Journal of Environment Management*, 2017, **186**: 192-200. <https://doi.org/10.1016/j.jenvman.2016.04.036>
- [15] Li N, Kang Y, Pan WJ, *et al.* Concentration and Transportation of Heavy Metals in Vegetables and Risk Assessment of Human Exposure to Bioaccessible Heavy Metals in Soil near a Waste-incinerator Site, South China. *Science of the Total Environment*, 2015, **521-522**: 144-151. <https://doi.org/10.1016/j.scitotenv.2015.03.081>
- [16] Cai LM, Xu ZC, Qi JY, *et al.* Assessment of Exposure to Heavy Metals and Health Risks among Residents near Tonglushan Mine in Hubei, China. *Chemosphere*, 2015, **127**: 127-135. <https://doi.org/10.1016/j.chemosphere.2015.01.027>
- [17] Ji K, Kim J, Lee M, *et al.* Assessment of Exposure to Heavy Metals and Health Risks among Residents near Abandoned Metal Mines in Goseong, Korea. *Environment Pollution*, 2013, **178**: 322-328. <https://doi.org/10.1016/j.envpol.2013.03.031>
- [18] Li B, Wang YH, Jiang Y, *et al.* The Accumulation and Health Risk of Heavy Metals in Vegetables around a Zinc Smelter in Northeastern China. *Environmental Science & Pollution Research International*, 2016, **23**(24): 1-13. <https://doi.org/10.1007/s11356-016-7342-5>
- [19] Eze VC, Onwukeme VI and Enyoh CE. Pollution status, ecological and human health risks of heavy metals in soil from some selected active dumpsites in Southeastern, Nigeria using energy dispersive X-ray spectrometer. *International Journal of Environmental Analytical Chemistry*, 2020. <https://doi.org/10.1080/03067319.2020.1772778>
- [20] Eze VC, Enyoh CE and Ndife CT. Soil Cationic Relationships, Structural and Fertility Assessment within selected active dumpsites in Nigeria. *Chemistry Africa*, 2020, **4**(4): 1-10. <https://doi.org/10.1007/s42250-020-00194-9>
- [21] APHA. (2005) Standard Methods for the Examination of Water and Wastewater 19 (American Public Health Association, APHA, AWWA. Washington. DC.
- [22] APHA. (1998) Standard methods for the Examination of Water and Wastewater (20th ed). New York: American Public Health, Association (APHA), American Water Works Association (AWWA), and Water Pollution Control Federation (WPCF).
- [23] Eze VC, Ndife CT and Muogbo MO. Carcinogenic and non-carcinogenic health risk assessment of heavy metals in Njaba River, Imo State, Nigeria. *Brazilian Journal of Analytical Chemistry*, 2021. <https://doi.org/10.30744/brjac.2179-3425.AR-05-2021>
- [24] Onwukeme VI and Eze VC. Identification of Heavy Metals Source within Selected Active Dumpsites in Southeastern Nigeria. *Environmental Analysis Health and Toxicology*, 2021, **36**(2): e2021008. <https://doi.org/10.5620/eaht.2021008>
- [25] Ibe FC, Chinedu NB and Ikpa CBC. Physicochemical analysis of some surface and groundwater sources in Etim Ekpo, LGA, Akwa Ibom State Nigeria. *Journal of Applied Science*, 2014, **17**(2): 11250-11260
- [26] Calo F and Parise M. Waste management and problems of groundwater pollution in karst environments in the context of a post-conflict scenario: the case of Mostar (Bosnia Herzegovina). *Habitat International*, 2009, **33**: 63. <https://doi.org/10.1016/j.habitatint.2008.05.001>
- [27] Wongsasuluk P, Chotpantararat S, Siriwong W, *et al.* Heavy metal contamination and human Health risk assessment in drinking water from shallow groundwater wells in an agricultural area in Ubong Ratchathani province, Thailand. *Environmental Geochemistry and Health*, 2014, **36**: 169-182. <https://doi.org/10.1007/s10653-013-9537-8>
- [28] Mishra S, Tiwary D, Ohri A, *et al.* Spatial analysis of groundwater quality around the MSW landfill site. *Nature, Environment and Pollution Technology*, 2018, **17**(3): 963-971.

- [29] Lele KC, Verla AW, Amaobi CE, *et al.* Health risks of consuming untreated borehole water from Uzoubi Umunna Orlu, Imo State Nigeria. *International Journal of Environmental Analytical Chemistry*, 2018, **5**(4): 1-7.
<https://doi.org/10.4172/2380-2391.1000250>
- [30] World Health Organization, WHO. (2003) *Environmental Health Criteria, 27, Guidelines on studies on Environmental Epidemiology*. Joint Sponsorship UNEP/ILO/WHO. Geneva.